

Genetic population structure of the stony coral *Seriatopora hystrix* in the Indo-Malay Archipelago



Vrije Universiteit Brussel

Hanneloor Heynderickx, Rosa Maria van der Ven, Puspita Sutrisno, Marc Kochzius
Marine Biology, Vrije Universiteit Brussel, Pleinlaan 2, 1050, Brussels, Belgium

Background

- Coral reefs are renowned to be some of the most diverse marine ecosystems on earth.
- Coral ecosystems contribute to local economies through tourism (diving & sedimentation) and fishing (the provision of food). More important they protect adjacent shorelines from wave action, storms and erosion [1].
- Approximately 75% of coral reefs worldwide are threatened by both local and global stressors with the highest threat in Southeast Asia where $\pm 95\%$ of the coral reefs are threatened [1-2].
- Marine protected areas (MPAs) can be a powerful management tool to increase connectivity and minimize coral resilience.

Research Objectives

- Acquiring knowledge on genetic population structure and hence connectivity patterns among subpopulations by measuring genetic diversity (allelic richness and heterozygosity).
- ❑ *Hypothesis: Allelic compositions will be different between the two studied sites and within Spermonde Archipelago, due to isolation by distance.*
- This study will provide necessary scientific information to support MPA implementations and effective management.

Species of Study

- ***Seriatopora hystrix*** is a hermaphroditic scleractinian brooding coral living in symbiosis with zooxanthellae.
- The species is native to East Africa, the Red Sea and western Indo-Pacific [3].
- Characterised by rapid growth, a variety of reproduction strategies and the capability for short- and long-distance dispersal [4].

Material & Methods

1. *Seriatopora hystrix* tissue samples: 303 samples from 13 locations (Fig. 1)
2. DNA extraction and quantification
3. Microsatellites selection and primer validation.
4. Multiplex PCR design
5. Fragment length analysis with a DNA capillary sequencer
6. Microsatellite scoring
7. Statistical analysis: Principal Components (PCoA), heterozygosity (F-statistics), GeneAIE (AMOVA + Mantel test), STRUCTURE (Bayesian cluster analysis)
8. Calculating genetic divergence within and among subpopulations




Fig. 1: Map of the Indo Malay Archipelago. [A] Pulau Seribu: 5 sampling site off Jakarta in Java. [B] Spermonde Archipelago: 8 sampling sites off the southwest coast of Sulawesi. ITF: Indonesian Throughflow, NECC: North Equatorial Counter Current and SEC: South Equatorial Current. Source (B): Landsat ETM + Satellite Image, Acquisition Year 2002.

Preliminary Results

10 Polymorphic microsatellites were selected
Two primermix setups were designed for multiplex PCR




Fig. 2: Two microsatellite primermix designs. The different microsatellites are color coded according to their fluorescent labels: 6FAM (blue), HEX (green), Cyanine 3 (yellow) and Atto 565 (red). Experimental lengths are indicated on top of the boxes, while theoretical lengths and repeat motifs (bold) are indicated in the grey boxes. The microsatellite primers were designed for species in the Red Sea (*) [6] and Australia [5]. SH: *Seriatopora hystrix*.

Acknowledgement

This research was funded by the German Federal Ministry of Education and Research (BMBF) in the framework of the SPICE project (Science for the Protection of Indonesian Coastal Marine Ecosystems). We would like to thank the institutions and individuals that have made our study possible: Janne Timm (Universität Bremen) for support during collection of samples; colleagues from Hasanuddin University (Indonesia) for logistical support during field work; the SPICE project is conducted and permitted under the governmental agreement between the BMBF and the Indonesian Ministry for Research and Technology (RISTEK), Indonesian Institute of Sciences (LIPI), Indonesian Ministry of Maritime Affairs and Fisheries (DKP), and Indonesian Agency for the Assessment and Application of Technology (BPPT). We thank the competent authorities for the permits.

References

1. Burke, L., Reynter, K., Spalding, M., Perry, A., 2011. Reefs at Risk Revisited. World Resources Institute, Washington, DC.
2. Wilkinson, C., 2008. Status of Coral Reefs of the World: 2008. Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, Townsville, Australia.
3. Borneman, E.H., 2001. Coral Reefs: Selection, Husbandry and Natural History. T.F.H. Publications, New Jersey, USA.
4. Stimson, J.S., 1978. Mode and timing of reproduction in some common hermatypic corals of Hawaii and Enewetak. Marine Biology 48, 173-184.
5. Underwood, J. N., Souter, P. B., Ballment, E. R., Lutz, A. H., & Van Oppen, M. J. H. (2006). Development of 10 polymorphic microsatellite markers from herbicide-bleached tissues of the brooding pocilloporid coral *Seriatopora hystrix*. Molecular Ecology Notes, 6, 176-178.
6. Maier, Elke, Ralph Tollrian, and Beate Nürnberg. "Development of species-specific markers in an organism with endosymbionts: microsatellites in the scleractinian coral *Seriatopora hystrix*." Molecular Ecology Notes 1.3 (2001): 157-159.

Hanneloor Heynderickx

MSc Marine and Lacustrine Science & Management
Marine Biology Research Group
Pleinlaan 2 - 1050 Brussels
Belgium

hanneloor.heynderickx@vub.ac.be

